This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.








FIG. 3




LANE


LANE

- 1. Cell culture supernatant
- 2. Heparin flow-through, 5% of load
- 3. Heparin flow-through, 38% of load
- 4. Heparin flow-through, 71% of load
- 5. Heparin flow-through, 100% of load
- 6. Heparin 500 mM NaCl eluate
- 7. Heparin 2M NaCl strip






- 1. Pierce Ab-Protein elution buffer.
- 2. 100 mM CAPS, pH 10.5.
- 3. 100 mM CAPS, 50% ethyl glycol, pH 10.5.
- 4. 100 mM Tris, 3.5 M, MgCl₂, 0.1% Tween 80, pH 8.



Elution Buffer pH



LANE

LANE

- 1. Cell culture supernatant
- 2. Heparin concentration eluate
- 3. MEP Hypercell eluate
- 4. Protein-A eluate